

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 340 783
A2

(12)

EUROPEAN PATENT APPLICATION

I D S

(21) Application number: 89108110.1

(51) Int. Cl. 4: C12M 3/00 , C12M 1/34

(22) Date of filing: 05.05.89

(30) Priority: 06.05.88 JP 109028/88
18.07.88 JP 176917/88

(43) Date of publication of application:
08.11.89 Bulletin 89/45

(84) Designated Contracting States:
CH DE FR GB LI

(71) Applicant: HITACHI, LTD.
6, Kanda Surugadai 4-chome Chiyoda-ku
Tokyo(JP)

(72) Inventor: Matsuzaki, Harumi
18-5, Daiharacho-2-chome
Hitachi-shi(JP)
Inventor: Haga, Ryoichi
11-3, Daiharacho-3-chome
Hitachi-shi(JP)
Inventor: Nishimura, Yuusaku
1-6, Kanesawacho-6-chome
Hitachi-shi(JP)

Inventor: Baba, Kenji
8-6, Mikanoharacho-1-chome
Hitachi-shi(JP)

Inventor: Maruhashi, Fumio
20-3, Ayukawacho-6-chome
Hitachi-shi(JP)

Inventor: Nishimura, Nobuko
5-1-407, Nagahoricho-2-chome
Katsuta-shi(JP)

Inventor: Ishida, Masahiko
20-8, Hanayamacho-1-chome
Hitachi-shi(JP)


Inventor: Saitoh, Setuo
26-3, Higashionumacho-2-chome
Hitachi-shi(JP)

(74) Representative: Ebbinghaus, Dieter et al
Patentanwälte v. Fünner, Ebbinghaus, Finck
Mariahilfplatz 2 & 3
D-8000 München 90(DE)

(54) Apparatus for culturing animal cells, method of culturing thereof and diagnostics of the culture.

(57) This invention provides an apparatus for culturing animal cells, which comprises diagnosis means to make it possible to diagnose by measuring sizes of the cells by means of image processing while a system of culturing the animal cells is cut off from the outside of the apparatus system, control means to control the culture conditions of a culturing cistern on the basis of a result of the diagnosis, and culture medium regenerating means to remove waste contents in a culture medium. In the apparatus of this invention, the culture system and diagnosis system are cut off from the outside of the apparatus system. Therefore, no infection of miscellaneous microorganism, etc., occurs, nor does consumption of valuable cells and products occur. Further, since it is possible to easily obtain data of the proportions of division potential-possessing cells and living cells, the environmental factors to the culturing cistern can be easily and instantly controlled based on the data, and therefore, the culturing efficiency can be enhanced.

FIG. 12



Xerox Copy Centre

EP 0 340 783 A2

APPARATUS FOR CULTURING ANIMAL CELLS, METHOD OF CULTURING THEREOF AND DIAGNOSTICS OF THE CULTURE

BACKGROUND OF THE INVENTION

FIELD OF THE INVENTION

5

This invention relates to the culturing of animal cells, and particularly it relates to an apparatus for culturing animal cells which makes it possible to measure and diagnose the activity status of animal cells under culturing while maintaining the physiological conditions of the animal cells, a culturing method and diagnostics of the culture.

10

THE RELATED ART STATEMENT

The production of new medical drugs by the culturing of animal cells gives as small an output as some
15 ng- μ g/ 10^6 cells \cdot day. The improvement of the drug productivity requires (1) high densification of cells and formation of a high capacity apparatus and (2) achievement of high secretory activity of cells. In order to attain the above (2), it is necessary to measure activities of cells in a culture medium (survival, division, secretion, etc.) under the condition that the physiological conditions of the cells are maintained and the system is closed. However, under the present situation, it is general practice to determine proportions of
20 living and dead cells according to a stain method in which a cell-staining agent is incorporated into a culture medium taken from a culturing cistern.

In the above stain method, a staining agent is incorporated, etc., i.e., the measurement cannot be made under the condition that the physiological conditions of cells are maintained, and it is therefore not possible to return the culture medium to the culturing cistern. Accordingly, the measurement system is under the
25 open system and it is impossible to prevent infection of miscellaneous microorganisms, etc. The rate of proliferation of animal cells is as small as 1/100 of that of microorganisms. Hence, the inclusion of one microorganism in the culture system means an end of new drug production activity.

On the other hand, with regard to cells being cultured in a microscopically observable container such as a microplate, dish or culturing bottle, there is a proposal of a method by which to determine the status of
30 cells, alive or dead, by image processing [Japanese Patent Kokai (laid-open) No. 201332/1987]. This method is applicable to species which is being cultured by using the above small containers. However, it is difficult to apply the above method to an apparatus of a practical culture scale.

Further, the above method is merely to determine the "living or dead" status of cells, and there is no industrial method to determine the activity status thereof at present.

35

OBJECT AND SUMMARY OF THE INVENTION

It is an object of this invention to provide a method of diagnosing the activity and culture status of cells
40 sampled for the diagnosis, while maintaining the physiological conditions of the cells, or a method of culturing cells which makes it possible to diagnose activity status of the cells without exerting any harmful influence on the culture system, and an apparatus for working the above method.

The present inventors have investigated animal cells in a culture medium and consequently found that the size of animal cells are grouped into the following three regions.

- 45
- (a) a region of only living cells (normal distribution)
 - (b) a region of both living and dead cells (homogeneous distribution)
 - (c) a region of only dead cells

Accordingly, it is possible to make a diagnosis on status of animal cells, which are being cultured, by
50 measuring the cell diameters of the animal cells.

The first aspect of this invention is directed to animal cell culturing apparatus which has a device for the above diagnosis. The embodiments thereof are as follows.

(1) An apparatus for culturing animal cells, which has a diagnosis device to make it possible to diagnose activity status of animal cells by measuring the sizes of the cells by means of image processing while maintaining their physiological conditions.

5 (2) An apparatus for culturing animal cells, which has, within a culture system of animal cells, a diagnosis device to make it possible to diagnose by measuring the sizes of the cells by means of image processing, and which is constituted to cut off the culture system from the outside the system.

(3) An apparatus for culturing animal cells, which has, within a cistern for culturing animal cells, a diagnosis device to make it possible to diagnose by measuring sizes of the cells by means of image processing.

10 (4) An apparatus for culturing animal cells, which has a system for extracting part of a cell culture medium from an animal cell culturing cistern, a diagnosing device to make it possible to diagnose by measuring sizes of the cells by means of image processing, and a system for returning measured culture cells and a medium of the cells to the culturing cistern have means to cut off themselves from the outside the individual systems.

15 (5) An apparatus for culturing animal cells, which has diagnosis means to make it possible to diagnose by measuring sizes of the cells by means of image processing while a system for culturing the animal cells is cut off from the outside the system, and control means to control the culture conditions of the culture cistern on the basis of the diagnostic result.

20 (6) An apparatus for culturing animal cells, which has a culturing cistern with control device to control environmental factors within the culturing cistern of animal cells, a diagnosis device to make it possible to diagnose the cells under culturing while maintaining physiological conditions of the cells, by measuring the sizes of the cell in a cell culture medium by means of image processing, circulation means to circulate the culture medium between the diagnosis device and the culturing cistern, and control means to control the control device of the culturing cistern on the basis of the result of image processing from the diagnosing device.

25 (7) An apparatus for culturing animal cells, which has diagnosing means to make it possible to diagnose by measuring sizes of the cells by means of image processing while a system for culturing the animal cells is cut off from the outside the apparatus system, control means to control the culture conditions a culturing cistern on the basis of a result of the diagnosis, and culture solution regenerating means to remove waste contents in a culture medium.

30 (8) An apparatus for culturing animal cells in any one of the above embodiments, which has a diagnosis device to diagnose activity status of the cells, under culturing, by measuring, by means of image processing, a distribution pattern of sizes of the cells in steady state of a culturing cistern of the cells and sizes of the cells at any cell-culturing time.

35 The second aspect of this invention relates to culturing method, and its embodiments are as follows.

(1) A method of culturing animal cells, which comprises taking part of a culture medium out of a culturing cistern, measuring sizes of cell particles in the culture solution by image processing while maintaining their physiological conditions, diagnosing activity status of the cells by a distribution of the sizes of the cell, and controlling the culture conditions of the culturing cistern on the basis of the diagnostic result.

40 (2) A method of culturing animal cells, which comprises, while a culture system is cut off from the outside the system, taking part of a culture medium out of a culturing cistern measuring sizes of cell in the culture medium by image processing, diagnosing activity status of the cells by a distribution of the sizes of the cell, and controlling the culture conditions of the culturing cistern on the basis of the diagnostic result.

45 (3) A method of culturing animal cells in any of the above embodiments (1) and (2), in which the culture solution used for the above diagnosis is cultured while recovering same into the culturing cistern.

The third aspect of this invention relates to diagnostics, and its embodiments are as follows.

50 (1) A method of diagnosing animal cells which comprises preliminarily diagnosing sizes of living cells and dead cells from a distribution pattern of sizes of cell in a culturing cistern of the animal cells in steady state, and measuring a distribution of the sizes of the cell at any culturing time by means of image processing and by using the above diagnostic result as a standard, thereby to diagnose activity status of animal cells.

55 (2) A method of diagnosing animal cells which comprises taking part of a culture medium out of a culturing cistern, changing a cells-suspending culture medium, which has been introduced into an observation portion cell while maintaining their physiological conditions, to image signals by image taking means, converting the analog signals of the image into digital signals by image processing means, storing brightnesses of pixels of the image in image memory means as a multi-value image data, calculating the

stored multi-value image data by cell image recognition means for a particle size distribution of all of the cells whose images are taken, and comparing the resultant distribution with the size distribution of the cells in culture steady state, thereby to diagnose activity status of the cells.

(3) A device for diagnosing animal cells, which has connection means to connect to a culturing cistern of the animal cells and means of measuring sizes of cell particles in a cell culture medium taken out of the culturing cistern through the above connecting means, by means of image processing while maintaining physiological conditions.

This invention can be applied to culturing animal cell strains such as human lymphoblast (IM-9: U.S. A.T.C.C.), rat liver cell JTC-1, mouse-mouse hybridoma (STK-1: U.S. A.T.C.C.), etc.

Needless to say that this invention is applicable to culturing of any cells that exhibit a cell size distribution as shown in Figure 1 which will be discussed later.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows experimental data for sizes and distribution states of animal cells.

Figure 2 shows a model for states of growth, division and death of animal cells.

Figures 3 to 5 are schematic views showing a device for diagnosing cell activity in this invention.

Figures 6 and 7 are schematic views showing the diagnosis of cell activity and control system of culturing cistern in this invention.

Figures 8 to 9 show typical views for a method of processing images of cells which are being cultured.

Figure 10 shows an example of the resultant cell diameter distribution.

Figure 11 shows an apparatus for diagnosing cell activity as one embodiment of this invention.

Figure 12 shows a device for controlling diagnosis of cell activity as one embodiment of this invention.

Figures 13 and 14 schematically show perspective views of a cell for diagnosing cells as one embodiment of this invention.

Figure 15 shows a constitution of a system for culturing animal cells.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

According to Figure 1, the judgement on the "living or dead" state of cells will be explained.

Figure 1 shows one example of cell size distributions obtained by microscopic observation of about 100 cells fixed by the stain method from rat liver cells JTC-1 in a steady state during a culturing process. In the Figure, that portion of the distribution which is indicated by hatching shows living cells, and that portion of the distribution which is indicated by void shows dead cells. The sizes of cells are distributed in the range from 10 μm and some to 40 μm and some. And the living cells are distributed on the smaller-number side and dead cells are distributed on the larger-number side. The above distribution consists of

- ① a region of only living cells,
- ② a region of both living and dead cells, and
- ③ a region of only dead cells.

On the basis of the above characteristic constitution of distribution, it is possible to determine proportions of living cells, dead cells and division potential-possessing cells. The living cells/dead cells ratio in the above region of both living and dead cells is 1:1, and the living cells in the region of both living and dead cells do not have any division potential. The living cells distributed in the above region ① have division potential.

The above distribution of cell sizes is in agreement with progressive phenomena of growth, division and death of cells typically shown in Figure 2. Statistically, when a cell grows to a certain size (two-dimensional size: its area is taken as 1), the cell undergoes division, and physically, its area after the division is 0.64 of the area which the cell had before the division. When reproduced cells grow to a size having an area of 1, the cells undergo division. The areas of division potential-possessing cells range between 0.64 and 1. A cell, which does not undergo division any longer, further grows over the area of 1 until its final death.

After a cell dies, cytoplasm such as mitochondria, etc., flow out of the cell, and the cell swells to increase its size apparently. Its nucleus disappears and finally the cell is dispersed in a culture medium. In

correspondence between the course of growth, division and death and the distribution of cell sizes shown in Figure 1, the state of area of 1 corresponds to a boundary between the distributions ① and ②, and the state of final death of the cell corresponds to a boundary between the distributions ② and ③. An actual course of cell culturing shows variations in the sizes of cells which finally die, and therefore the distribution ② appears. In the living cells/dead cells ratio in the distribution ②, the smaller the cell sizes are, the larger the proportion of living cells is, and the larger the cell sizes are, the smaller the proportion of living cells is. The living cells/dead cells ratio in the distribution ② is nearly 1:1.

As mentioned above, it is possible to determine the living/dead ratio and division activities of cells by measuring cell size distributions. The measurement of cell size distributions can be carried out without incorporating any foreign substance into a culture system. Therefore, the above activities of cells can be diagnosed without isolating the cells from their culture system.

EXAMPLES

Figure 3 shows a device for diagnosing cells as one embodiment of this invention. The apparatus comprises a culturing cistern 1, a culture medium observing portion 2, means 3 of introducing a culture medium to the observing portion 2, a measuring device 4 to measure sizes of cells in the observing portion 2, an analyzer 8 to determine a proportion of division potential-possessing cells (or proportion of division potential-not possessing cells) or a proportion of living cells (or proportion of dead cells) on the basis of a cell size distribution, and means 6 to return a culture medium in the observing portion 2 to a culture medium in the culturing cistern. The means 3 and 6 and the observing portion 2 are constituted such that the culture medium can be recycled under a condition of its isolation from the outside the culture system.

The culture medium in the culturing cistern is partially fed to the observing portion 2, and cells in the partially fed culture medium in the observing portion 2 are measured by the measuring device 4 for a cell size distribution. An equivalent diameter or area may be used to represent the cell size. When the measurement is finished, the culture medium in the observing portion 2 is returned into the culturing cistern 1. An information on the cell size is transmitted from the measuring device 4 to the analyzer 8, and the analyzer 8 determines activity of cells (division activity, life-maintenance activity) on the basis of the above information to display the result.

In general, the culture medium is intermittently fed from the culturing cistern 1 to the measuring portion 2. However, there are some types of measuring devices which can measure the above distribution while the culture medium is fed continuously.

Figure 4 shows a device for diagnosing cell activity as one embodiment of this invention, in which an image processing device is used in the above measuring device 4. The image processing device comprises an image taking means 4' and an image processing means 5. An image of cells is transmitted from the image taking means 4' to the image processing means 5. Preferably, the image taking means 4' has a magnifying function such as a microscopic function. The flow rate of a culture medium in the observing portion 2 may be about a rate of precipitation of cells in a stationary culture medium.

Figure 5 shows detailed constitutions of the image processing device 5 and the analyzer 8, and their actions will be explained hereinbelow.

At first, image signals obtained by the image taking device 4' are transmitted to an A/D converter 51 of the image processing device 5. The A/D converter 51 converts analog signals of an image to digital signals. That is, the image is not only divided vertically and horizontally into small pixels, but also A/D converted image signals are formed into a multi-element image in which the pixels have digital brightness values. For example, an image is divided vertically and horizontally into 256 by 256 pixels and digitized with resolution of 128-step brightnesses. The A/D converted image signals are stored in an image memory 51M. The multi-element image stored in the image memory 51M is transmitted to a cell image recognition circuit 60. The cell image recognition circuit 60 recognizes all of the cells regardless of their life or death, calculates their number and sizes, and calculates their particle diameter distribution. The calculation results are transmitted to the analyzer 8, and the analyzer calculates, on the basis of the above calculation results, the numbers of living cells and dead cells and the number of division potential-possessing cells, and at the same time diagnoses activity status of the grouped cells on the basis of these calculation results.

Figure 6 shows an embodiment of multi-value image data stored in the image memory 51M. The multi-element image stands for an image of which the pixels are different in the brightness, as explained previously. Some cells have large sizes and the other cells have small sizes. In Figure 6, cells having small particle diameters represent living cells, and cells having large particle diameters represent dead cells. Cells having diameters in between represent those in the state in which living cells and dead ones are co-present.

The image in Figure 6 is transmitted to the cell image recognition circuit 60, which recognizes all of the cells regardless of their life or death, and calculates the number, size, form and brightness of the cells, etc. This Example will explain a calculation of the size and number.

A pre-processing circuit 61 effects the pre-processing in order to recognize the cells from the multi-element image of the image memory 51M. Examples of the preprocessing include smoothing, brightness emphasis, outline emphasis, etc. However, in order to recognize cells, mere spatial filtering treatment is not sufficient in accuracy, and particularly, brightness emphasizing treatment before the spatial filtering treatment is effective. The maximum brightness emphasizing treatment is one type of nonlinear neighbourhood calculation, and its specific details will be explained according to Figure 7. In a region of 3 x 3 pixels (of which the brightnesses are indicated by k_1 to k_9) having a pixels of brightness k_5 in the middle as shown in Figure 7, the maximum brightness in the region is newly substituted in the picture element of brightness k_5 . The calculation formula is as follows.

$$k_5 = \text{MAX} [k_1 \sim k_9] \quad (1)$$

For example, if k_1 is the highest brightness among k_1 to k_9 , i.e., $\text{MAX} [k_1 \sim k_9] = k_1$, then the brightness of the pixel having a brightness k_5 is changed to k_1 . All the picture elements are treated in the same way as mentioned here to obtain a new multi-value image data. This treatment has a function to emphasize the brightness of the central portion of a cell since the central portion has high brightness.

Further, there is also a function to erase dark portions of a cell outline. In addition, in the pre-processing circuit 61, it is further effective to carry out spatial filtering treatment in order to emphasize bright portions by using a mask pattern shown in Figure 8. In Figure 8, p's represent positive values, and m's represent negative values. This treatment makes it possible to selectively emphasize cells alone.

A binalization circuit 62 achieves binalization in which portions having higher brightnesses than the given brightness k_a are set as "1" and the other portions as "0". Figure 9 shows an image obtained by this binalization of the image of Figure 6. In Figure 9, portions indicated by slanting lines show background, and void portions show cells. If cells are large and interiors of the cells are binalized as blank, blank-filling treatment is carried out, although its explanation is omitted here. By the above procedures, it is possible to selectively recognize cells alone. The parameters, m, p, k_a , etc., which are necessary for the preprocessing and binalization, are set by a conditioning circuit 66.

With regard to an image binalized by the binalization circuit 62, a labeling circuit 63 labels numbers on cells therein and counts their numbers, and every cell is subjected to the following treatment. That is, a next area calculation circuit 64 calculates an area of every cell. Then, a circuit 5, for calculation of a cell diameter distribution, calculates a diameter by classifying areas of the cells. For example, the classification is effected every 1 μm . In this case, a circle-equivalent diameter, etc., may be used as a typical cell diameter. The calculated cell diameter distribution is transmitted to the analyzer 8.

The analyzer 8 receives values of the cell size distribution and diagnoses activities of the cell groups totally. An example of the resultant cell diameter distribution is shown in Figure 10. This principle is accomplished by forming a model of a relationship between growth of cells (increase in cell diameter) and their diameter distribution. That is, in the analyzer 8, a living cell calculation circuit 81A, a dead cell calculation circuit 81B and a co-present cell calculation circuit 81C calculate numbers of grouped cells, respectively. In other words, a formula of living cells (2), formula of dead cells (3) and formula of living and dead cells co-present state (4) are formed into models with separate distributions of each diameters. Namely, provided that cell diameters present in boundaries of the grouped cells are d_{am} and d_{me} as shown in Figure 10:

$$\text{living cell } d \leq d_{am} : N(d_a) = F_a(d) \quad (2)$$

$$\text{dead cell } d > d_{me} : N(d_e) = F_e(d) \quad (3)$$

$$\text{co-presence } d_{am} < d \leq d_{me} : N(d_m) = F_m(d) \quad (4)$$

$$\text{total number of cells : } N_t(d) = N(d_a) + N(d_e) + N(d_m) \quad (5)$$

In the above formulae, d_a , d_e and d_m represent cell diameters of living cells only, dead cells only and living and dead cells co-present state, respectively, and $N(d_a)$, $N(d_e)$ and $N(d_m)$ represent numbers of such cells, respectively. Further, $F_a(d)$, $F_e(d)$ and $F_m(d)$ are functions of cell diameter distributions, and $F_a(d)$ is a normal distribution, $F_m(d)$ is a homogeneous distribution and $F_e(d)$ is a monotone decrease type distribution (e.g., exponential distribution). As mentioned here, the living cell calculation circuit 81A, dead cell calculation circuit 81B and co-present cell calculation circuit 81C carry out calculations to apply models-forming particle diameter distributions to the cell particle diameter distribution transmitted from the cell image recognition circuit 60, thereby to determine the cell diameters d_{am} and d_{me} which are boundaries of the individual distributions. If the boundary cell diameters d_{am} and d_{me} are determined, $N(d_a)$, $N(d_e)$ and $N(d_m)$ can be easily obtained from the cell diameter distribution transmitted from the cell image recognition circuit 60. The above $N(d_a)$, $N(d_e)$ and $N(d_m)$ can also be obtained by inputting d_{am} and d_{me} through the

conditioning circuit 82 in the steady state.

The number of cells in each of the distributions can be calculated as mentioned above. Therefore, various activities of the cell groups as a whole can be diagnosed. Diagnostics will be discussed concretely hereinbelow.

5 An activity calculation circuit 83 calculates, for example, a cell survival rate ϵ in the following formula.

$$\epsilon = \frac{N(d_a) + \frac{1}{2}(d_m)}{N_t(d)} \quad \dots\dots (6)$$

10

In the activity calculation circuit 83, a division potential-possessing cell ratio η is calculated in the following formula.

15

$$\eta = \frac{N(d_a)}{N(d_a) + \frac{1}{2}N(d_m)} \quad \dots\dots (7)$$

20

Similarly, in the activity calculation circuit 83, a dead cell ratio R_e is calculated in the following formula.

$$R_e = \frac{N(d_e) + \frac{1}{2}N(d_m)}{N_t(d)} \quad \dots\dots (8)$$

25

30 In addition to the above, in the activity calculation circuit 83, temporal changes of living cells $\{N(d_a) + 1/2N(d_m)\}$ and dead cells $\{N(d_e) + 1/2N(d_m)\}$ are calculated, i.e., a rate of proliferation and a rate of death are calculated in the following formulae.

$$35 \quad \text{Rate of proliferation} = \frac{d\{N(d_a) + \frac{1}{2}N(d_m)\}}{dt} \quad \dots (9)$$

$$40 \quad \text{Rate of death} = \frac{d\{N(d_e) + \frac{1}{2}N(d_m)\}}{dt} \quad \dots\dots (10)$$

45 On the basis of these formulae, cells are diagnosed for proliferation, activity decrease or death state. For example, even if R_e is large, the higher rate of proliferation indicates the higher proliferation potency and activity of the cells. Even if R_e is small, the higher rate of death indicates not only rapid death of the cells but also low activity of the cells. The activities of cell groups are diagnosed as mentioned above.

50 These diagnostic results are inputted to an inference engine 95. Also inputted to the inference engine 95 are values measured by a temperature sensor, a pH sensor, dissolved oxygen (DO) sensor and a dissolved carbon dioxide gas (DCO₂) sensor as cell culture environmental factors. On the other hand, assorted experimental and theoretical data of proliferation and metabolism of cells are stored in a data base 90. The inference engine 95 infers status of cells on the basis of information of the data base, measured values of culture environment and diagnostic results of image-measurement of cell groups, and inputs a command to a computer as necessary in order to control the culture environment. For example, when DO decreases to lower the proliferation rate of cells, the inference engine 95 infers DO shortage and increases the rate of bubble generation by a compressor to increase DO. When pH lowers to increase the rate of death, the inference engine 95 infers that pH is too low, and increases pH by feeding an alkali agent, adjusting the partial pressure of carbon dioxide gas, or the like.

55

Figure 11 shows an apparatus for diagnosing cell activity as one embodiment of this invention. This apparatus is an integral constitution of the culturing cistern 1 and observing portion 2 shown in Figure 4, in which a culture medium in the culturing cistern is partially fed to the observing portion 2 by a spinner flow in the culturing cistern and the culture medium is returned from the observing portion 2 into the culturing

5 cistern.

Figure 12 shows a device for controlling diagnosis of cell activity as one embodiment of this invention. This apparatus comprises a culturing cistern 1 having means (1', 1'', 1''', etc.) of adjusting environmental factors to a culture medium within the culturing cistern (nutrition contents concentration, dissolved oxygen concentration, concentration of waste products such as ammonia, lactic acid or the like, ages of cells, product concentration, etc.), an observing portion 2, an introduction device 3 to introduce a culture medium to the observing portion 2, a measuring device 4 to measure a distribution of cell sizes in the observing portion 2, an analyzer 8 to determine, from the cell size distribution, a proportion of division potential-possessing cells (or proportion of division potential-not possessing cells) or a proportion of living cells (or proportion of dead cells), and means 6 of returning the culture medium in the observing portion 2 into the culture medium within the culturing cistern. The above means 3 and 6 and the observing portion 2 are constituted such that the culture medium is recycled under the condition that it is cut off from the outside the culture system.

The culture medium in the culturing cistern is partially fed to the observing portion 2, and cells in the fed culture medium are measured by the measuring device 4 for a distribution of their sizes. Nominal diameters, areas, etc., may be used to represent sizes of the cells. The culture medium used for measurement in the observing portion 2 is returned into the culturing cistern 1. An information of a cell size distribution is transmitted from the measuring device 4 to the analyzer 8, which determines cell activities (division activity, survival activity) on the basis of the above information. Further, the analyzer 8 properly adjusts the environmental conditions to the culture medium in the culturing cistern by controlling the above adjusting means 1', 1'', 1''', etc., on the basis of the resultant cell activities.

25 The method of controlling the environmental conditions to the culture medium in the culturing cistern will be explained hereinbelow.

By using a proportion η of division potential-possessing cells (= number of division potential-possessing cells/number of living cells) and a proportion ϵ of living cells (= number of living cells/number of total cells), the number N of living cells to be found in t hours' time can be represented by the following formula.

$$35 \quad N = \frac{1}{2} N_0 \left\{ \frac{1 - (2\eta)^{\frac{t}{T} + 1}}{1 - (2\eta)} + 1 \right\} \dots\dots (11)$$

wherein if $(n + 1)T > t \geq nT$, then $\frac{t}{T} = n$

40 In the above formula, N_0 represents a total cell number when $t=0$ and T is a cell cycle, and n is 0 or a positive integer.

That is, it is possible to simulate the number of living cells in t hours' time by using formula (11). On the basis of the result of the above simulation, it is also possible to control the environmental conditions to the culture medium in the culturing cistern. In addition to the dissolved oxygen concentration, nutrition contents concentration and waste contents concentration in the culture medium, the above environmental conditions include the following values P and Q .

$$P = \text{cell concentration} \times \text{stay time} \quad (12)$$

$$50 \quad Q = \frac{\text{cell concentration} \times \text{stay time}}{\text{buoyant substance concentration in fed nutrition content liquid}} \dots\dots (13)$$

55 By controlling the above value P or Q such that it is in a given range while considering the above environmental conditions, it is possible to control various activities of cells (respiratory activity, proliferation activity, secretory activity, endogenous respiratory activity, etc.). That is, it is possible to control activities optionally; for example, the proliferation activity can be enhanced more than the other activities, the

secretory activity can be enhanced more than the other activities, and the like.

The following is an explanation of a measuring device cell to measure the cell sizes in this invention.

Figure 13 is a perspective view showing one embodiment of said cell. Figure 14 is a typical perspective view of a cell portion in which a culture medium of cells flows.

5 The observing portion cell 2 has a cell structure as shown in Figure 14, which is formed by using, as a spacer 11, a film of polytetraethylene (having a thickness of about 100 μm) with a hexagonal window, and placing the film between two glass sheets 12 and 13. The two glass sheets are fastened by two metal supporter frames 16 and 17 and screws (not shown). The cell is provided with an inflow tube 14 of a cell culture medium and an outflow tube 15 thereof. In addition, the observing portion cell can be subjected to
10 sterilization treatment under heat (120° C).

Further, the above cell is placed on a stage 18 which is movable freely in the direction of X-Y, and is provided with a constitution which makes it possible to make observation through the measuring device 4 placed above the cell and by light emitted from a light source (not shown) placed in the lower portion of the stage.

15 A culture medium is introduced into the cell by the inflow tube 14 from a circulation pipe (not shown) connected to the culturing cistern, and the stage and an observing device are manipulated to observe sizes of animal cells. In diagnosis of the activity status of cells, an image of cells, which is taken by a image making device (indicated as 4' in Figure 4) attached to the measuring device 4, is processed by an image processing means (indicated as 5 in Figure 4), and the resultant data is diagnosed by an analyzing means
20 (indicated as 8 in Figure 4).

The culture medium containing the measured cells is returned to the culturing cistern through the outflow tube 15.

As mentioned above, since this embodiment makes it possible to diagnose the activity status of cells in the state where the culture system and diagnosis system are cut off from the outside the apparatus system,
25 there is not any risk of miscellaneous microorganisms infection, nor is there any loss of products such as a culture medium, etc.

As discussed hereinabove, in this invention, the culture system and diagnosis system are cut off from the outside the apparatus system. Therefore, there is no infection of miscellaneous microorganisms, etc., and valuable cells and products are not consumed. Further, since it is possible to easily obtain data of the
30 proportion of division potential-possessing cells and the proportion of living cells, it is easily and instantly control the environmental factors to the culturing cistern on the basis of the data. Therefore, the culturing efficiency can be enhanced.

One embodiment of the animal cell culture system, which is precedent to the foregoing diagnosis of animal cell activities, will be explained hereinbelow.

35 Figure 15 shows a constitution of one embodiment of a cell culture system usable in this invention.

In a culturing cistern 102, cells are cultured by being dispersed and suspended in a culture medium 101. The culturing cistern is provided, in its bottom portion, with a nozzle to aerate the medium with an oxygen-containing gas, through which oxygen necessary for growth of cells and carbon dioxide gas necessary for pH adjustment are fed. That is, air from an air compressor 160 and an oxygen-containing gas
40 from a container 170 are flow-controlled such that the dissolved oxygen concentration is constant, and further a carbon dioxide gas from a container 180 is flow-controlled such that the pH is between 7.0 and 7.6. These gases are passed through a sterilizing filter 107 and then blown in through the nozzle 104. The gas blown into the solution starts the culture medium flowing by forming bubbles and ascending toward the surface of the medium. A draft tube 103 is provided in order to achieve the effective flow of the culture
45 medium started by the blown gas. Bubbles having reached the medium surface form a bubble layer without breakage, and they are therefore broken effectively by bringing them into contact with a antifoaming agent layer 105 formed of a water repellent material above the medium surface.

Gas is discharged, as discharge gas 124, out of the culturing cistern through a mist separator 106 and a filter 107. In addition, the culturing cistern 102 is provided with various devices such as means of
50 introducing culture medium to the observing portion, means of adjusting a dissolved gas concentration and pH to constant levels, means of sterilizing the interiors of the cistern and tubes, means of charging and discharging culture medium, means of maintaining temperature at a given level, sensors to detect a dissolved oxygen concentration, pH, temperature, internal pressure of the cistern, medium surface level of the cistern, etc., and the like. However, these devices are not shown in Figure 15.

55 The culturing cistern 102 is provided with a filtration device 108 in its lower portion. The filtration device 108 is provided with a porous hollow fiber membrane filter 110 inside and with a filter auxiliary 109 on or around the filter. The filter auxiliary 109 is supported by a support filter 111.

It is to be noted that the hollow fiber membrane filter 110 changes its form or exposes itself from the

layer of the filter auxiliary 109 when the solution moves rapidly at the time of sterilization, backwash, etc. However, its form is not specially limited.

The culture solution exchange is carried out as follows. In Figure 15, valves 131 and 132 are closed, a valve 133 is opened and a pump 122 is actuated to introduce a culture medium 101 from the culturing cistern 102 to the filtration device 108. Cells in the culture medium are collected on the surfaces of the filter auxiliary 109 and porous hollow fiber membrane filter 110 when the culture solution passes therethrough. The porous hollow fiber membrane filter 110 is selected such that it has an opening degree of preventing the passage of particles of the filter auxiliary 109 and the cells. A filtrate of the culture medium, from which the cells are filtered, is stored in a culture medium storage tank 115.

After a given amount of the culture medium is discharged from the culturing cistern 102, the pump 122 is stopped and the valve 133 is closed. Then, the valves 131 and 132 are opened, and pumps 119 and 120 are actuated. The pumps 119 and 120 feed a fresh culture medium from a fresh culture medium storage tank 112 to the filtration device. In this case, the pump 119 feeds the fresh culture medium from the bottom portion of the filtration device 108, and its feed rate is set so as to float and spread the filler layer of the filter auxiliary 109. The pump 120 reversely charges a fresh culture medium under pressure onto the porous hollow fiber membrane filter 110 for its backwash, thereby to separate solid substances such as cells, etc., from the surface of the filter 110. In addition, the flow rates of the pumps 119 and 120 are adjusted so as not to cause dissipation of the filter auxiliary 109 from around the filtration device.

The cells collected on the filtration auxiliary 109 and the hollow fiber membrane filter 110 in the above filtration step were released out of the filler layer by the spreading of the filtration auxiliary 109. The specific gravity, size and form of the particles of the filtration auxiliary 109 are selected such that the settling rate of the particles is higher than that of the cells. Therefore, the cells are floated up to the upper portion of the spreaded filler layer, and back to the culturing cistern 102 together with the fresh culture medium. In addition, numeral 121 indicates a pump and numeral 125 indicates a tubing system from the culture filtrate tank.

Claims

1. An apparatus for culturing animal cells, which comprises a diagnosis device to make it possible to diagnose activity status of animal cells by measuring the sizes of the cells by means of image processing while maintaining their physiological conditions.

2. An apparatus for culturing animal cells, which has, within a culture system of animal cells, a diagnosis device to make it possible to diagnose by measuring the sizes of the cells by means of image processing, and which is constituted to cut off the culture system from the outside of the system.

3. An apparatus for culturing animal cells, which has, within a cistern for culturing animal cells, a diagnosis device to make it possible to diagnose by measuring sizes of the cells by means of image processing.

4. An apparatus for culturing animal cells, wherein a system of extracting part of a cell culture medium from an animal cell culturing cistern, a diagnosis device to make it possible to diagnose by measuring sizes of the cells by means of image processing, and a system of returning measured culture cells and a suspension of the cells to the culturing cistern have means of cutting off themselves from the outside of the individual systems.

5. An apparatus for culturing animal cells, which comprises diagnosis means to make it possible to diagnose by measuring sizes of the cells by means of image processing while a system of culturing the animal cells is cut off from the outside of the system, and control means to control the culture conditions of the culture cistern on the basis of the diagnostic result.

6. An apparatus for culturing animal cells, which comprises a culturing cistern with a control device to control environmental factors to the culturing cistern of the animal cells, a diagnosis device to make it possible to diagnose the cells under culturing by measuring the sizes of the cell particles in a cell culture medium by means of image processing while maintaining physiological conditions of the cells, circulation means to circulate the culture medium between the diagnosis device and the culturing cistern, and control means to control the control device of the culturing cistern on the basis of the result of image processing from the diagnosis device.

7. An apparatus for culturing animal cells, which comprises diagnosis means to make it possible to diagnose by measuring sizes of the cells by means of image processing while a system of culturing the animal cells is cut off from the outside of the apparatus system, control means to control the culture conditions in a culturing cistern on the basis of a result of the diagnosis, and culture solution regenerating means to remove waste contents in a culture solution.

8. An apparatus for culturing animal cells according to any one of claims 1 to 7, which has a diagnosis device to diagnose activity status of the cells under culturing by measuring, by means of image processing, a distribution pattern of cell sizes in steady state of a culturing cistern of the cells and sizes of the cells at any cell-culturing time.

9. A method of culturing animal cells, which comprises taking part of a culture medium out of a culturing cistern, measuring sizes of cell in the culture solution by image processing while maintaining their physiological conditions, diagnosing activity status of the cells from a distribution of the sizes of the cell, and controlling the culture conditions of the culturing cistern on the basis of the diagnostic result.

10. A method of culturing animal cells, which comprises, while their culture system is cut off from the outside of the system, taking part of a culture medium out of a culturing cistern, measuring sizes of cell in the culture medium by image processing, diagnosing activity status of the cells by a distribution of the sizes of the cell, and controlling the culture conditions of the culturing cistern on the basis of the diagnostic result.

11. A method of culturing animal cells according to any of claims 9 and 10, wherein the culture medium used for the above diagnosis is cultured while recovering same into the culturing cistern.

12. A method of diagnosing animal cells which comprises preliminarily diagnosing sizes of living cells and dead cells from a distribution pattern of sizes of cell in a culturing cistern of the animal cells in steady state, and measuring a distribution of the sizes of the cell at any culture time by means of image processing and by using the above diagnostic result as a standard, thereby to diagnose activity status of the animal cells.

13. A method of diagnosing animal cells which comprises taking part of a culture medium out of a culturing cistern, changing a cells-containing culture medium, which has been introduced into an observation portion cell while maintaining their physiological conditions, to image signals by image taking means, converting the analog signals of the image into digital signals by image processing means, storing brightnesses of pixels of the image in image memory means as a multi-element image, calculating the stored multi-element image by cell image recognizing means for a particle size distribution of all of the cells which are picked up, and comparing the resultant distribution with the cell size distribution in culture steady state, thereby to diagnose activity status of the cells.

14. A device for diagnosing animal cells, which comprises connection means to connect to a culturing cistern of the animal cells and means of measuring sizes of cell in a cell culture medium taken out of the culturing cistern through the above connection means, by means of image processing while maintaining their physiological conditions.

15. A method of culturing animal cells which comprises discharging part of a culture medium with a filter membrane and feeding a fresh culture medium while carrying out a liquid culture of the animal cells, the filter membrane being a hydrophobic filter membrane and being backwashed intermittently with a fresh culture medium.

16. An apparatus for liquid-culturing living organism cells which comprises means of feeding a gas useful for the cells which are being cultured, means of feeding a culture medium and discharge means of discharging a waste culture medium, said discharge means being attached through culture medium-filtration means of a hydrophobic filter membrane, said hydrophobic filter membrane having means thereby to achieve its backwash with a culture medium.

50

55

FIG. 1




FIG. 2




FIG. 3




FIG. 4





FIG. 5



0340783A2

FIG. 6




FIG. 7

k_1	k_2	k_3
k_4	k_5	k_6
k_7	k_8	k_9

FIG. 8

m	m	m	m	m	m
m	P	P	P	P	m
m	P	P	P	P	m
m	P	P	P	P	m
m	P	P	P	P	m
m	m	m	m	m	m

FIG. 9




FIG. 10




FIG. 11




FIG. 12






FIG. 13





FIG. 14







FIG. 15



12

EUROPEAN PATENT APPLICATION

21 Application number: 89108110.1

51 Int. Cl.⁵: C12M 3/00, C12M 1/34,
C12M 1/36

22 Date of filing: 05.05.89

30 Priority: 06.05.88 JP 109028/88
18.07.88 JP 176917/88

43 Date of publication of application:
08.11.89 Bulletin 89/45

84 Designated Contracting States:
CH DE FR GB LI

88 Date of deferred publication of the search report:
18.07.90 Bulletin 90/29

71 Applicant: HITACHI, LTD.
6, Kanda Surugadai 4-chome
Chiyoda-ku, Tokyo 100(JP)

72 Inventor: Matsuzaki, Harumi
18-5, Daiharacho-2-chome
Hitachi-shi(JP)
Inventor: Haga, Ryoichi
11-3, Daiharacho-3-chome
Hitachi-shi(JP)
Inventor: Nishimura, Yuusaku

1-6, Kanesawacho-6-chome
Hitachi-shi(JP)

Inventor: Baba, Kenji

8-6, Mikanoharacho-1-chome
Hitachi-shi(JP)

Inventor: Maruhashi, Fumio

20-3, Ayukawacho-6-chome
Hitachi-shi(JP)

Inventor: Nishimura, Nobuko

5-1-407, Nagahoricho-2-chome
Katsuta-shi(JP)

Inventor: Ishida, Masahiko

20-8, Hanayamacho-1-chome
Hitachi-shi(JP)

Inventor: Saitoh, Setuo

26-3, Higashionumacho-2-chome
Hitachi-shi(JP)


74 Representative: Ebbinghaus, Dieter et al
Patentanwälte v. Fünér, Ebbinghaus, Finck
Mariahilfplatz 2 & 3
D-8000 München 90(DE)

54 Apparatus for culturing animal cells, method of culturing thereof and diagnostics of the culture.

57 This invention provides an apparatus for culturing animal cells, which comprises diagnosis means to make it possible to diagnose by measuring sizes of the cells by means of image processing while a system of culturing the animal cells is cut off from the outside of the apparatus system, control means to control the culture conditions of a culturing cistern on the basis of a result of the diagnosis, and culture medium regenerating means to remove waste contents in a culture medium. In the apparatus of this invention, the culture system and diagnosis system are cut off from the outside of the apparatus system. Therefore, no infection of miscellaneous microorganism, etc., occurs, nor does consumption of valuable cells and products occur. Further, since it is possible to easily obtain data of the proportions of division

potential-possessing cells and living cells, the environmental factors to the culturing cistern can be easily and instantly controlled based on the data, and therefore, the culturing efficiency can be enhanced.

FIG. 12





European Patent
Office

EUROPEAN SEARCH REPORT

Application number

EP 89 10 8110

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Y	DE-A-3 139 310 (VSESOJUZYNYJ NAUCNO-ISSLEDOVATEL'SKIY INSTITUT BIOSINTEZA BELKOVYCH VESCESTV) * Page 12, paragraph 3 - page 33, paragraph 4, lines 1-9; figures 1-7 *	1-14	C 12 M 3/00 C 12 M 1/34 C 12 M 1/36
Y	DE-A-3 516 529 (GESELLSCHAFT FÜR STRAHLEN- UND UMWELTFORSCHUNG mbH) * Page 5, paragraph 3 - page 11, paragraph 3; claims 1-6; figures 1,2 *	1-14	
D,A	PATENT ABSTRACTS OF JAPAN; & JP-A-62 201 332 (SUMITOMO ELECTRIC IND. LTD) 05-09-1987	1,13	TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
A	PATENT ABSTRACTS OF JAPAN, vol. 9, no. 153 (C-288)(1876), 27th June 1985; & JP-A-60 31 888 (HITACHI SEISA- KUSHO K.K.)18-02-1985	1,13	C 12 M
E	EP-A-0 336 974 (SUMITOMO ELECTRICAL IND. LTD) * Column 6, line 15 - column 7, line 14; column 12, lines 3-36; column 15, line 1 - column 16, ./.	1-3,5, 7-10, 12-14	
FURTHER INFORMATION			
Place of search		Date of completion of the search	Examiner
The Hague		05-01-1990	GROENENDIJK
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO Form 1503 03 82



CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid.
- namely claims:
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

X LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions.

namely:

1. Claims 1-14: Method and apparatus for culturing animal cells comprising diagnosis of activity status of cells and feed-back control
2. Claims 15-16: Method and apparatus for culturing animal cells comprising use of hydrophobic filter membrane

- ☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid.
- namely claims:
- ☒ None of the further search fees has been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims.

namely claims: 1-14



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 3)
	line 32; column 16, line 49 - column 17, line 5; column 17, line 48 - column 18, line 41; claims 1,12,16; figure 1 * -----		
			TECHNICAL FIELDS SEARCHED (Int. Cl. 3)
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
<div><div>CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</div><div>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</div></div>			

EPO Form 1503 03 82